arXiv:1303.4826v1 [math.NT] 20 Mar 2013

CONGRUENCES FOR THE k DOTS BRACELET PARTITION FUNCTIONS

SUPING CUI AND NANCY SHANSHAN GU

Abstract. By finding the congruent relations between the generating function of the 5 dots bracelet partitions and that of the 5-regular partitions, we get some new congruences modulo 2 for the 5 dots bracelet partition function. Moreover, for a given prime p, we study the arithmetic properties modulo p of the k dots bracelet partitions.

1. Introduction

In [1], Andrews and Paule studied the broken k-diamond partitions by using MacMahon's partition analysis, and gave the generating function of $\Delta_k(n)$ which denotes the number of the broken k-diamond partitions of n:

$$\sum_{n=0}^{\infty}\Delta_k(n)q^n=\frac{(-q;q)_\infty}{(q;q)_\infty^2(-q^{2k+1};q^{2k+1})_\infty}. \tag{1.1}$$

In [1], They proved the following arithmetic theorem for $\Delta_1(n)$.

Theorem 1.1. *[1, Theorem 5] For $n \geq 0$,*

$$\Delta_1(2n+1)\equiv 0 \pmod 3.$$

Meanwhile, they posed some conjectures related to $\Delta_2(n)$. For other study of the arithmetic of the broken k-diamond partitions, see [3, 7, 9–12, 14, 17]. In [4], Fu found a combinatorial proof of Theorem 1.1 for $\Delta_1(n)$, and introduced a generalization of the broken k-diamond partitions which he called the k dots bracelet partitions. The number of this kind of partitions of n is denoted by $\mathfrak{B}_k(n)$, and the generating function of $\mathfrak{B}_k(n)$ is stated as follows.

$$\sum_{n=0}^{\infty}\mathfrak{B}_k(n)q^n=\frac{(-q;q)_\infty}{(q;q)_\infty^{k-1}(-q^k;q^k)_\infty},\ k\geq 3.$$

In [4], Fu proved the following congruences for $\mathfrak{B}_k(n)$.

Theorem 1.2. *[4, Theorem 3.3] For $n > 0$, $k \geq 3$, if $k = p^r$ is a prime power, we have*

$$\mathfrak{B}_k(2n+1)\equiv 0 \pmod p.$$

Date: March 21, 2013.

1991 *Mathematics Subject Classification.* 11P83, 05A17.

Key words and phrases. partitions, broken k-diamond partitions, k dots bracelet partitions, congruences.

Theorem 1.3. *[4, Theorem 3.5] For any $k \geq 3$, s an integer between 1 and $p-1$ such that $12s+1$ is a quadratic nonresidue modulo p, and any $n \geq 0$, if $p \mid k$ for some prime $p \geq 5$ say $k = pm$, then we have*

$$\mathfrak{B}_k(pn+s) \equiv 0 \pmod{p}.$$

Theorem 1.4. *[4, Theorem 3.6] For $n \geq 0$, $k \geq 3$ even, say $k = 2^m l$, where l is odd, we have*

$$\mathfrak{B}_k(2n+1) \equiv 0 \ (mod\ 2^m).$$

Later, in [13], Radu and Sellers found some new congruences for $\mathfrak{B}_k(n)$.

Theorem 1.5. *[13, Theorem 1.4] For all $n \geq 0$,*

$$\begin{gathered}\mathfrak{B}_5(10n+7) \equiv 0 \ (mod\ 5^2),\\ \mathfrak{B}_7(14n+11) \equiv 0 \ (mod\ 7^2), \ \ and\\ \mathfrak{B}_{11}(22n+21) \equiv 0 \ (mod\ 11^2).\end{gathered}$$

In this paper, we continue to study the arithmetic of the k dots bracelet partitions. First, we recall two kinds of partitions which are used in this paper.

A partition of a positive integer n is a nonincreasing sequence of positive integers whose sum is n. Let $p(n)$ denote the number of partitions of n. We know that

$$\sum_{n=0}^{\infty} p(n)q^n = \frac{1}{(q;q)_\infty}. \tag{1.2}$$

If ℓ is a positive integer, then a partition is called ℓ-regular partition if there is no part divisible by ℓ. Let $b_\ell(n)$ denote the number of ℓ-regular partitions of n. The generating function of $b_\ell(n)$ is stated as follows.

$$\sum_{n=0}^{\infty} b_\ell(n)q^n = \frac{(q^\ell;q^\ell)_\infty}{(q;q)_\infty}. \tag{1.3}$$

In section 2, based on an identity given by Ramanujan in [16] and a congruence for the generating function of $b_5(2n)$ given by Hirschhorn and Sellers in [8], we obtain two congruences modulo 2 for $B_5(n)$. Meanwhile, by finding a congruent relation between the generating function of $\mathfrak{B}_5(n)$ and that of $b_5(n)$, we get many infinite family of congruences modulo 2 for $\mathfrak{B}_5(n)$. In section 3, for a given prime p, by means of a p-dissection identity of $f(-q)$ given by the authors in [6] and three classical congruences for $p(n)$ given by Ramanujan in [15,16], we get more congruences modulo p for $\mathfrak{B}_k(n)$.

In the following, we list some definitions and identities which are frequently used in this paper.

As usual, we follow the notation and terminology in [5]. For $|q| < 1$, the q-shifted factorial is defined by

$$(a;q)_\infty = \prod_{k=0}^{\infty}(1-aq^k) \quad \text{and} \quad (a;q)_n = \frac{(a;q)_\infty}{(aq^n;q)_\infty}, \text{ for } n \in \mathbb{C}.$$

The Legendre symbol is a function of of a and p defined as follows:

$$\left(\frac{a}{p}\right)=\begin{cases}1, & \text{if } a \text{ is a quadratic residue modulo } p \text{ and } a\not\equiv 0 \pmod{p},\\ -1, & \text{if } a \text{ is a quadratic non-residue modulo } p,\\ 0, & \text{if } a\equiv 0 \pmod{p}.\end{cases}$$

Jacobi's triple product identity [2, Theorem 1.3.3]: for $z\neq 0$ and $|q|<1$,

$$\sum_{n=-\infty}^{\infty} z^n q^{n^2}=(-zq,-q/z,q^2;q^2)_\infty. \tag{1.4}$$

Ramanujan's general theta function $f(a,b)$ is defined by

$$f(a,b):=\sum_{n=-\infty}^{\infty} a^{\frac{n(n+1)}{2}} b^{\frac{n(n-1)}{2}}, \qquad |ab|<1.$$

A special case of $f(a,b)$ is stated as follows.

$$f(-q):=f(-q,-q^2)=\sum_{n=-\infty}^{\infty}(-1)^n q^{\frac{n(3n-1)}{2}}=(q;q)_\infty.$$

2. Congruences modulo 2 for $\mathfrak{B}_5(n)$

First, we recall an identity given by Ramanujan in [16, p. 212].

$$(q;q)_\infty=\frac{(q^{10},q^{15},q^{25};q^{25})_\infty}{(q^5,q^{20};q^{25})_\infty}-q(q^{25};q^{25})_\infty-q^2\frac{(q^5,q^{20},q^{25};q^{25})_\infty}{(q^{10},q^{15};q^{25})_\infty}. \tag{2.1}$$

For convenience, we set

$$a(q)=\frac{(q^{10},q^{15};q^{25})_\infty}{(q^5,q^{20};q^{25})_\infty} \qquad \text{and} \qquad b(q)=\frac{(q^5,q^{20};q^{25})_\infty}{(q^{10},q^{15};q^{25})_\infty}=\frac{1}{a(q)}.$$

Then, we can rewrite (2.1) as

$$(q;q)_\infty=(q^{25};q^{25})_\infty(a(q)-q-q^2b(q)). \tag{2.2}$$

In [8], Hirschhorn and Sellers obtained the following congruence for the generating function of $b_5(2n)$.

$$\sum_{n=0}^{\infty} b_5(2n)q^n \equiv (q^2;q^2)_\infty \pmod{2}. \tag{2.3}$$

By means of (2.2) and (2.3), we have the following results for $\mathfrak{B}_5(n)$.

Theorem 2.1. *For $n\geq 0$, we have*

$$\mathfrak{B}_5(10n+6)\equiv 0 \pmod{2},$$
$$\mathfrak{B}_5(10n+8)\equiv 0 \pmod{2}.$$

Proof. First, we have

$$\begin{aligned}\sum_{n=0}^{\infty}\mathfrak{B}_5(n)q^n &= \frac{(-q;q)_\infty}{(q;q)_\infty^4(-q^5;q^5)_\infty}\\ &= \frac{(q^2;q^2)_\infty(q^5;q^5)_\infty}{(q;q)_\infty^5(q^{10};q^{10})_\infty}\\ &\equiv \frac{(q^2;q^2)_\infty(q^5;q^5)_\infty}{(q^4;q^4)_\infty(q^{10};q^{10})_\infty(q;q)_\infty} \pmod 2\\ &\equiv \frac{1}{(q^2;q^2)_\infty(q^{10};q^{10})_\infty}\frac{(q^5;q^5)_\infty}{(q;q)_\infty} \pmod 2\\ &= \frac{1}{(q^2;q^2)_\infty(q^{10};q^{10})_\infty}\cdot\sum_{n=0}^{\infty}b_5(n)q^n.\end{aligned}$$

Then,

$$\begin{aligned}\sum_{n=0}^{\infty}\mathfrak{B}_5(2n)q^n &\equiv \frac{1}{(q;q)_\infty(q^5;q^5)_\infty}\cdot\sum_{n=0}^{\infty}b_5(2n)q^n \pmod 2\\ &\equiv \frac{(q^2;q^2)_\infty}{(q;q)_\infty(q^5;q^5)_\infty} \pmod 2 \qquad \text{by (2.3)}\\ &\equiv \frac{(q;q)_\infty}{(q^5;q^5)_\infty} \pmod 2.\end{aligned}$$

According to (2.2), we have

$$\sum_{n=0}^{\infty}\mathfrak{B}_5(2n)q^n \equiv \frac{(q^{25};q^{25})_\infty}{(q^5;q^5)_\infty}(a(q)-q-q^2b(q)) \pmod 2. \tag{2.4}$$

Therefore, we get

$$\begin{aligned}\mathfrak{B}_5(2(5n+3)) &= \mathfrak{B}_5(10n+6)\equiv 0 \pmod 2,\\ \mathfrak{B}_5(2(5n+4)) &= \mathfrak{B}_5(10n+8)\equiv 0 \pmod 2.\end{aligned}$$

□

Lemma 2.2. *We have*

$$\sum_{n=0}^{\infty}\mathfrak{B}_5(10n+2)q^n \equiv \sum_{n=0}^{\infty}b_5(n)q^n \pmod 2.$$

Proof. Due to (2.4), we get

$$\sum_{n=0}^{\infty}\mathfrak{B}_5(2(5n+1))q^n = \sum_{n=0}^{\infty}\mathfrak{B}_5(10n+2)q^n \equiv \frac{(q^5;q^5)_\infty}{(q;q)_\infty} = \sum_{n=0}^{\infty}b_5(n)q^n \pmod 2.$$

□

In [6], the authors found many infinite family of congruences modulo 2 for $b_5(n)$.

Theorem 2.3. *[6, Theorem 3.17] For any prime $p \geq 5$, $\left(\frac{-10}{p}\right) = -1$, $\alpha \geq 1$, and $n \geq 0$, we have*

$$b_5(4 \cdot p^{2\alpha} n + \frac{(24i + 7p)p^{2\alpha-1} - 1}{6}) \equiv 0 \pmod{2},$$

where $i = 1, 2, \ldots, p-1$.

Theorem 2.4. *[6, Theorem 3.20] For $\alpha \geq 0$ and $n \geq 0$, we have*

$$\begin{aligned} b_5(4 \cdot 5^{2\alpha+1} n + \frac{31 \cdot 5^{2\alpha} - 1}{6}) &\equiv 0 \pmod{2}, \\ b_5(4 \cdot 5^{2\alpha+1} n + \frac{79 \cdot 5^{2\alpha} - 1}{6}) &\equiv 0 \pmod{2}, \\ b_5(4 \cdot 5^{2\alpha+2} n + \frac{83 \cdot 5^{2\alpha+1} - 1}{6}) &\equiv 0 \pmod{2}, \\ b_5(4 \cdot 5^{2\alpha+2} n + \frac{107 \cdot 5^{2\alpha+1} - 1}{6}) &\equiv 0 \pmod{2}. \end{aligned}$$

Therefore, combining Lemma 2.2 with Theorem 2.3 and Theorem 2.4, we obtain some more congruences for $\mathfrak{B}_5(n)$.

Theorem 2.5. *For any prime $p \geq 5$, $\left(\frac{-10}{p}\right) = -1$, $\alpha \geq 1$, and $n \geq 0$, we have*

$$\mathfrak{B}_5(40 \cdot p^{2\alpha} n + \frac{5 \cdot (24i + 7p)p^{2\alpha-1} + 1}{3}) \equiv 0 \pmod{2},$$

where $i = 1, 2, \ldots, p-1$.

For example, by setting $p = 17$, $i = 6$, and $\alpha = 1$ in Theorem 2.5, we have the following congruence.

$$\mathfrak{B}_5(11560n + 7452) \equiv 0 \pmod{2}.$$

Theorem 2.6. *For $\alpha \geq 1$ and $n \geq 0$, we have*

$$\begin{aligned} \mathfrak{B}_5(8 \cdot 5^{2\alpha} n + \frac{31 \cdot 5^{2\alpha-1} + 1}{3}) &\equiv 0 \pmod{2}, \\ \mathfrak{B}_5(8 \cdot 5^{2\alpha} n + \frac{79 \cdot 5^{2\alpha-1} + 1}{3}) &\equiv 0 \pmod{2}, \\ \mathfrak{B}_5(8 \cdot 5^{2\alpha+1} n + \frac{83 \cdot 5^{2\alpha} + 1}{3}) &\equiv 0 \pmod{2}, \\ \mathfrak{B}_5(8 \cdot 5^{2\alpha+1} n + \frac{107 \cdot 5^{2\alpha} + 1}{3}) &\equiv 0 \pmod{2}. \end{aligned}$$

3. CONGRUENCES MODULO p FOR $\mathfrak{B}_k(n)$

In [6], the authors studied a p-dissection identity of $f(-q)$ for a given prime $p \geq 5$.

Theorem 3.1. *[6, Theorem 2.2] For any prime $p \geq 5$, we have*

$$f(-q) = \sum_{\substack{k=-\frac{p-1}{2} \\ k \neq \frac{\pm p-1}{6}}}^{\frac{p-1}{2}} (-1)^k q^{\frac{3k^2+k}{2}} f(-q^{\frac{3p^2+(6k+1)p}{2}}, -q^{\frac{3p^2-(6k+1)p}{2}}) + (-1)^{\frac{\pm p-1}{6}} q^{\frac{p^2-1}{24}} f(-q^{p^2}),$$

where $\pm$ depends on the condition that $(\pm p-1)/6$ should be an integer. Meanwhile, we claim that $(3k^2+k)/2$ and $(p^2-1)/24$ are not in the same residue class modulo p for $-(p-1)/2 \leq k \leq (p-1)/2$ and $k \neq (\pm p-1)/6$.

According to the above theorem, we have the following result.

Lemma 3.2. *For any prime $p \geq 5, n \geq 0$, and $r \geq 1$, if $k = p^r$ is a prime power, then for $1 \leq \alpha \leq (r+1)/2$, we have*

$$\sum_{n=0}^{\infty} \mathfrak{B}_k(p^{2\alpha-1}n + \frac{p^{2\alpha}-1}{12})q^n \equiv \left((-1)^{\frac{\pm p-1}{6}}\right)^{\alpha} \frac{(q^{2p};q^{2p})_\infty}{(q^{2p^{r-(2\alpha-1)}};q^{2p^{r-(2\alpha-1)}})_\infty} \pmod{p},$$

where $\pm$ depends on the condition that $(\pm p-1)/6$ should be an integer.

Proof. We prove the lemma by induction on α. For $k = p^r$, in [4], Fu stated the following fact

$$\begin{aligned}\sum_{n=0}^{\infty} \mathfrak{B}_k(n)q^n &= \frac{(-q;q)_\infty}{(q;q)_\infty^{k-1}(-q^k;q^k)_\infty} \\ &= \frac{(q^2;q^2)_\infty}{(q;q)_\infty^{k}(-q^k;q^k)_\infty} \\ &\equiv \frac{(q^2;q^2)_\infty}{(q^k;q^k)_\infty(-q^k;q^k)_\infty} \pmod{p} \\ &= \frac{(q^2;q^2)_\infty}{(q^{2k};q^{2k})_\infty} \pmod{p}.\end{aligned}$$

Due to Theorem 3.1, for any prime $p \geq 5$, we have

$$\sum_{n=0}^{\infty} \mathfrak{B}_k(pn + \frac{p^2-1}{12})q^n \equiv (-1)^{\frac{\pm p-1}{6}} \frac{(q^{2p};q^{2p})_\infty}{(q^{2p^{r-1}};q^{2p^{r-1}})_\infty} \pmod{p}.$$

That means the lemma holds for $\alpha = 1$. Suppose that lemma holds for α. Now we prove the case for $\alpha + 1$. For

$$\sum_{n=0}^{\infty} \mathfrak{B}_k(p^{2\alpha-1}n + \frac{p^{2\alpha}-1}{12})q^n \equiv \left((-1)^{\frac{\pm p-1}{6}}\right)^{\alpha} \frac{(q^{2p};q^{2p})_\infty}{(q^{2p^{r-(2\alpha-1)}};q^{2p^{r-(2\alpha-1)}})_\infty} \pmod{p}.$$

Then

$$\begin{aligned}\sum_{n=0}^{\infty} \mathfrak{B}_k(p^{2\alpha-1}(pn) + \frac{p^{2\alpha}-1}{12})q^n &= \sum_{n=0}^{\infty} \mathfrak{B}_k(p^{2\alpha}n + \frac{p^{2\alpha}-1}{12})q^n \\ &\equiv \left((-1)^{\frac{\pm p-1}{6}}\right)^{\alpha} \frac{(q^2;q^2)_\infty}{(q^{2p^{r-2\alpha}};q^{2p^{r-2\alpha}})_\infty} \pmod{p}. \quad (3.1)\end{aligned}$$

Using Theorem 3.1 again, we have

$$\begin{aligned}&\sum_{n=0}^{\infty}\mathfrak{B}_k(p^{2\alpha}(pn+\frac{p^2-1}{12})+\frac{p^{2\alpha}-1}{12})q^n\\ =&\sum_{n=0}^{\infty}\mathfrak{B}_k(p^{2\alpha+1}n+\frac{p^{2\alpha+2}-1}{12})q^n\\ \equiv&\left((-1)^{\frac{\pm p-1}{6}}\right)^{\alpha+1}\frac{(q^{2p};q^{2p})_\infty}{(q^{2p^{r-(2\alpha+1)}};q^{2p^{r-(2\alpha+1)}})_\infty}\pmod{p}.\end{aligned}$$

Therefore the lemma holds for $\alpha+1$. □

According to Lemma 3.2, we have the following results.

Theorem 3.3. *For any prime $p\geq 5, n\geq 0$, and $r\geq 1$, if $k=p^r$ is a prime power, then we have the following two cases:*

(1) *For $i=1,2,\cdots,p-1$ and $1\leq\alpha\leq r/2$, we have*

$$\mathfrak{B}_k(p^{2\alpha}n+\frac{(12i+p)p^{2\alpha-1}-1}{12})\equiv 0\pmod{p}.$$

(2) *Let j be an integer between 1 and $p-1$ and $12j+1$ is a quadratic nonresidue modulo p. For $n\geq 0$ and $1\leq\alpha\leq(r-1)/2$, we have*

$$\mathfrak{B}_k(p^{2\alpha+1}n+\frac{(12j+1)p^{2\alpha}-1}{12})\equiv 0\pmod{p}.$$

Proof. According to Lemma 3.2, when $1\leq\alpha\leq r/2$, for $i=1,2,\cdots,p-1$, we have

$$\mathfrak{B}_k(p^{2\alpha-1}(pn+i)+\frac{p^{2\alpha}-1}{12})=\mathfrak{B}_k(p^{2\alpha}n+\frac{(12i+p)p^{2\alpha-1}-1}{12})\equiv 0\pmod{p}.$$

For $1\leq\alpha\leq(r-1)/2$, according to (3.1) and Theorem 3.1, we know that the powers of q modulo p congruent to $2\cdot(3k^2+k)/2$ for $-(p-1)/2\leq k\leq(p-1)/2$ in the expansion of $(q^2;q^2)_\infty$. So we have

$$\begin{gathered}j\equiv 2\cdot\frac{3k^2+k}{2}\pmod{p},\\ 12j+1\equiv(6k+1)^2\pmod{p}.\end{gathered}$$

Therefore, if $12j+1$ is a quadratic nonresidue modulo p, then we have

$$\sum_{n=0}^{\infty}\mathfrak{B}_k(p^{2\alpha}(pn+j)+\frac{p^{2\alpha}-1}{12})q^n\equiv 0\pmod{p}.$$

□

Based on Lemma 3.2 and the generating functions of $p(n)$ and $b_\ell(n)$, we get the following congruent relations.

Theorem 3.4. *For any prime $p \geq 5$, $\alpha \geq 1$, and $n \geq 0$, if $k = p^{2\alpha-1}$ is a prime power, then we have*

$$\sum_{n=0}^{\infty} \mathfrak{B}_k(2p^{2\alpha-1}n + \frac{p^{2\alpha}-1}{12})q^n \equiv \left((-1)^{\frac{\pm p-1}{6}}\right)^{\alpha} \sum_{n=0}^{\infty} b_p(n)q^n \pmod{p}, \tag{3.2}$$

$$\sum_{n=0}^{\infty} \mathfrak{B}_k(2p^{2\alpha-1}n + \frac{p^{2\alpha}-1}{12})q^n \equiv \left((-1)^{\frac{\pm p-1}{6}}\right)^{\alpha} (q^p;q^p)_\infty \sum_{n=0}^{\infty} p(n)q^n \pmod{p}. \tag{3.3}$$

Proof. Set $r = 2\alpha - 1$ in Lemma 3.2. Then $k = p^{2\alpha-1}$. So we get

$$\sum_{n=0}^{\infty} \mathfrak{B}_k(p^{2\alpha-1}n + \frac{p^{2\alpha}-1}{12})q^n \equiv \left((-1)^{\frac{\pm p-1}{6}}\right)^{\alpha} \frac{(q^{2p};q^{2p})_\infty}{(q^2;q^2)_\infty} \pmod{p}.$$

Therefore,

$$\begin{aligned}\sum_{n=0}^{\infty} \mathfrak{B}_k(p^{2\alpha-1}(2n) + \frac{p^{2\alpha}-1}{12})q^n &= \sum_{n=0}^{\infty} \mathfrak{B}_k(2p^{2\alpha-1}n + \frac{p^{2\alpha}-1}{12})q^n \\ &\equiv \left((-1)^{\frac{\pm p-1}{6}}\right)^{\alpha} \frac{(q^p;q^p)_\infty}{(q;q)_\infty} \pmod{p}.\end{aligned}$$

□

Combining (3.3) with the three famous congruences for $p(n)$ given by Ramanujan in [15, 16]

$$p(5n+4) \equiv 0 \pmod{5}, \tag{3.4}$$

$$p(7n+5) \equiv 0 \pmod{7}, \tag{3.5}$$

$$p(11n+6) \equiv 0 \pmod{11}, \tag{3.6}$$

we get the following results.

Corollary 3.5. *For $\alpha \geq 1$ and $n \geq 0$, we have*

$$\begin{aligned}\mathfrak{B}_{5^{2\alpha-1}}(2\cdot 5^{2\alpha}n + \frac{101\cdot 5^{2\alpha-1}-1}{12}) &\equiv 0 \pmod{5},\\ \mathfrak{B}_{7^{2\alpha-1}}(2\cdot 7^{2\alpha}n + \frac{127\cdot 7^{2\alpha-1}-1}{12}) &\equiv 0 \pmod{7},\\ \mathfrak{B}_{11^{2\alpha-1}}(2\cdot 11^{2\alpha}n + \frac{155\cdot 11^{2\alpha-1}-1}{12}) &\equiv 0 \pmod{11}.\end{aligned}$$

Proof. According to (3.3), we have

$$\begin{aligned}\sum_{n=0}^{\infty} \mathfrak{B}_{5^{2\alpha-1}}(2\cdot 5^{2\alpha-1}n + \frac{5^{2\alpha}-1}{12})q^n &\equiv (-1)^{\alpha}(q^5;q^5)_\infty \sum_{n=0}^{\infty} p(n)q^n \pmod{5},\\ \sum_{n=0}^{\infty} \mathfrak{B}_{7^{2\alpha-1}}(2\cdot 7^{2\alpha-1}n + \frac{7^{2\alpha}-1}{12})q^n &\equiv (-1)^{\alpha}(q^7;q^7)_\infty \sum_{n=0}^{\infty} p(n)q^n \pmod{7},\\ \sum_{n=0}^{\infty} \mathfrak{B}_{11^{2\alpha-1}}(2\cdot 11^{2\alpha-1}n + \frac{11^{2\alpha}-1}{12})q^n &\equiv (q^{11};q^{11})_\infty \sum_{n=0}^{\infty} p(n)q^n \pmod{11}.\end{aligned}$$

Based on (3.4), (3.5), and (3.6), we get

$$\begin{aligned}\mathfrak{B}_{5^{2\alpha-1}}(2\cdot 5^{2\alpha-1}(5n+4)+\frac{5^{2\alpha}-1}{12}) &\equiv 0 \pmod 5,\\ \mathfrak{B}_{7^{2\alpha-1}}(2\cdot 7^{2\alpha-1}(7n+5)+\frac{7^{2\alpha}-1}{12}) &\equiv 0 \pmod 7,\\ \mathfrak{B}_{11^{2\alpha-1}}(2\cdot 11^{2\alpha-1}(11n+6)+\frac{11^{2\alpha}-1}{12}) &\equiv 0 \pmod{11}.\end{aligned}$$

□

Another congruence modulo p for $\mathfrak{B}_k(n)$ can be directly obtained from Lemma 3.2.

Theorem 3.6. *For any prime $p \geq 5$, $\alpha \geq 1$, and $n \geq 1$, if $k = p^{2\alpha}$ is a prime power, then we have*

$$\mathfrak{B}_k(p^{2\alpha-1}n+\frac{p^{2\alpha}-1}{12}) \equiv 0 \quad (mod\ p).$$

Proof. Set $r = 2\alpha$ in Lemma 3.2. Then $k = p^{2\alpha}$. So we have

$$\sum_{n=0}^{\infty}\mathfrak{B}_k(p^{2\alpha-1}n+\frac{p^{2\alpha}-1}{12})q^n \equiv \left((-1)^{\frac{\pm p-1}{6}}\right)^{\alpha} \pmod p.$$

□

Acknowledgements: This work was supported by the National Natural Science Foundation of China and the PCSIRT Project of the Ministry of Education.

References

[1] G. E. Andrews and P. Paule, MacMahon's partition analysis XI: Broken diamonds and modular forms, Acta Arith. 126 (2007), 281–294.

[2] B. C. Berndt, Number Theory in the Spirit of Ramanujan, American Mathematical Society, Providence, 2004.

[3] S. H. Chan, Some congruences for Andrews-Paule's broken 2-diamond partitions, Discrete Math. 308 (2008), 5735–5741.

[4] S. Fu, Combinatorical proof of one congruence for the broken 1-diamond partition and a generalization, Int. J. Number Theory 7 (2011), 133–144.

[5] G. Gasper and M. Rahman, Basic Hypergeometric Series, Second Ed., Cambridge University Press, Cambridge, 2004.

[6] S. P. Cui and N. S. S. Gu, Arithmetic properties of the ℓ-regular partitions, arXiv:1302.3693 [math.CO].

[7] M. D. Hirschhorn and J. A. Sellers, On recent congruence results of Andrews and Paule, Bull. Austral. Math. Soc. 75 (2007), 121–126.

[8] M. D. Hirschhorn and J. A. Sellers, Elementary proofs of parity results for 5-regular partitions, Bull. Austral. Math. Soc. 81 (2010), 58–63.

[9] M. Jameson, Congruences for broken k-diamond partitions, Ann. Comb., to appear.

[10] E. Mortenson, On the broken 1-diamond partition, Int. J. Number Theory 4 (2008), 199–218.

[11] P. Paule and S. Radu, Infinite families of strange partition congruences for broken 2-diamonds, Ramanujan J. 23 (2010), 409–416.

[12] S. Radu and J. A. Sellers, Parity results for broken k-diamond partitions and $(2k+1)$-cores, Acta Arith. 146 (2011), 43–52.

[13] S. Radu and J. A. Sellers, Congruences modulo squares of primes for Fu's k dots bracelet partitions, Int. J. Number Theory, to appear.
[14] S. Radu and J. A. Sellers, Infinite many congruences for broken 2-diamond partitions modulo 3, J. Comb. Number Theory, to appear.
[15] S. Ramanujan, Some properties of $p(n)$, the number of partitions of n, Proc. Cambridge Philos. Soc. 19 (1919), 210–213.
[16] S. Ramanujan, Collected Papers, Cambridge University Press, Cambridge, 1927; reprinted by Chelsea, New York, 1962; reprinted by the American Mathematical Society, Providence, RI, 2000.
[17] X. H. Xiong, Two congruences involving Andrews-Paule's broken 3-diamond partitions and 5-diamond partitions, Proc. Japan Acad. Ser. A Math. Sci. 87 (2011), 65–68.

(S. P. Cui) Center for Combinatorics, LPMC-TJKLC, Nankai University, Tianjin 300071, P. R. China

E-mail address: `jiayoucui@163.com`

(N. S. S. Gu) Center for Combinatorics, LPMC-TJKLC, Nankai University, Tianjin 300071, P. R. China

E-mail address: `gu@nankai.edu.cn`